

Table 7.82b  
Updated

Native Hawaiian Men Aged 40 and Over in Hawai'i Who Had a Digital Rectal Exam (DRE) and Prostate Specific Antigen (PSA) Test in Hawai'i: 2005-2012

| Year | Native Hawaiian Men Aged 40+             |             |                           |             |   |             |                                |             |
|------|--|-------------|---------------------------|-------------|---|-------------|--------------------------------|-------------|
|      | Percentage Who Had a Digital Rectal Exam |             |                           |             | Percentage Who Had a Prostate Specific Antigen Test |             |                                |             |
|      | Ever had a DRE                           |             | Last DRE within Past Year |             | Ever had a PSA Test                                 |             | Last PSA Test within Last Year |             |
|      | %  | C.I. (95%)  | %                         | C.I. (95%)  | %   | C.I. (95%)  | %                              | C.I. (95%)  |
| 2012 | (X)                                      |             | (X)                       |             | 29.4  | (21.0-39.5) | 9.3                            | ( 6.0-14.1) |
| 2011 | (X)                                      |             | (X)                       |             | 35.1  | (27.0-44.1) | 20.2                           | (13.8-28.6) |
| 2010 | 58.0                                     | (48.3-67.2) | 22.7                      | (16.4-30.6) | 49.0  | (39.9-58.2) | 30.2                           | (22.6-39.2) |
| 2009 | 53.4                                     | (40.3-66.0) | 15.6                      | ( 7.9-28.2) | 37.2  | (25.7-50.4) | 25.6                           | (15.3-39.5) |
| 2008 | 53.7                                     | (44.2-62.9) | 28.1                      | (20.1-37.7) | 40.2  | (31.6-49.5) | 31.7                           | (23.3-41.5) |
| 2007 | 65.2                                     | (56.4-73.0) | 28.5                      | (21.2-37.1) | 46.1  | (37.2-55.2) | 27.6                           | (20.8-35.7) |
| 2006 | 49.3                                     | (40.4-58.2) | 17.4                      | (12.2-24.2) | 28.0  | (21.3-36.0) | 19.7                           | (14.0-27.1) |
| 2005 | 66.0                                     | (49.7-79.3) | 24.8                      | (15.0-38.1) | 32.7  | (21.3-46.5) | 21.7                           | (13.0-34.1) |

(X) = Not reported/Question not asked

Source: Hawai'i State, Department of Health, Behavioral Risk Factor Surveillance System.